



# Assignment

## Human eye

- Near and far points of human eye are  
(a) 25 cm and infinite      (b) 50 cm and 100 cm      (c) 25 cm and 50 cm      (d) 0 cm and 25 cm
- A defective eye cannot see close objects clearly because their image is formed  
(a) On the eye lens      (b) Between eye lens and retina  
(c) On the retina      (d) Beyond retina
- Retina of eye acts like ..... of camera  
(a) Shutter      (b) Film      (c) Lens      (d) None of these
- A person who can see things most clearly at a distance of 10 cm. Requires spectacles to enable him to see clearly things at a distance of 30 cm. What should be the focal length of the spectacles  
(a) 15 cm (concave)      (b) 15 cm (convex)      (c) 10 cm      (d) 0
- An astronaut is looking down on earth's surface from a space shuttle at an altitude of 400 km. Assuming that the astronaut's pupil diameter is 5 mm and the wavelength of visible light is 500 nm. The astronaut will be able to resolve linear object of the size of about  
(a) 0.5 m      (b) 5 m      (c) 50 m      (d) 500 m
- A person uses a lens of power + 3D to normalise vision. Near point of hypermetropic eye is  
(a) 1 m      (b) 1.66 m      (c) 2 m      (d) 0.66 m
- The separation between two microscopic particles is measured  $P_A$  and  $P_B$  by two different lights of wavelength 2000 Å and 3000 Å respectively, then  
(a)  $P_A > P_B$       (b)  $P_A < P_B$       (c)  $P_A < 3/2 P_B$       (d)  $P_A = P_B$
- To remove myopia (short sightedness) a lens of power 0.66 D is required. The distant point of the eye is approximately  
(a) 100 cm      (b) 150 cm      (c) 50 cm      (d) 25 cm
- A person suffering from 'presbyopia' should use  
(a) A concave lens      (b) A convex lens  
(c) A bifocal lens whose lower portion is convex      (d) A bifocal lens whose upper portion is convex

10. The resolving limit of healthy eye is about  
 (a) 1' (b) 1" (c) 1° (d)  $\frac{1}{60}$  "
11. A person uses spectacles of power + 2D. He is suffering from  
 (a) Short sightedness or myopia (b) Long sightedness or hypermetropia  
 (c) Presbyopia (d) Astigmatism
12. The hyper metropia is a  
 (a) Short-side defect (b) Long-side defect  
 (c) Bad vision due to old age (d) None of these
13. A man cannot see clearly the objects beyond a distance of 20 cm from his eyes. To see distant objects clearly he must use which kind of lenses and of what focal length  
 (a) 100 cm convex (b) 100 cm concave (c) 20 cm convex (d) 20 cm concave
14. An eye specialist prescribes spectacles having a combination of convex lens of focal length 40 cm in contact with a concave lens of focal length 25 cm. The power of this lens combination in diopters is [IIT 1997 Cancelled]  
 (a) + 1.5 (b) -1.5 (c) +6.67 (d) -6.67
15. Two parallel pillars are 11 km away from an observer. The minimum distance between the pillars so that they can be seen separately will be  
 (a) 3.2 m (b) 20.8 m (c) 91.5 m (d) 183 m
16. A person cannot see objects clearly beyond 2.0 m. The power of lens required to correct his vision will be  
 (a) + 2.0 D (b) - 1.0 D (c) + 1.0 D (d) - 0.5 D
17. When objects at different distances are seen by the eye, which of the following remains constant  
 (a) The focal length of the eye lens (b) The object distance from the eye lens  
 (c) The radii of curvature of the eye lens (d) The image distance from the eye lens
18. A person wears glasses of power -2.0 D. The defect of the eye and the far point of the person without the glasses will be  
 (a) Nearsighted, 50 cm (b) Farsighted, 50 cm (c) Nearsighted, 250 cm (d) Astigmatism, 50 cm
19. A person is suffering from the defect astigmatism. Its main reason is  
 (a) Distance of the eye lens from retina is increased (b) Distance of the eye lens from retina is decreased  
 (c) The cornea is not spherical (d) Power of accommodation of the eye is decreased
20. Myopia is due to  
 (a) Elongation of eye ball (b) Irregular change in focal length  
 (c) Shortening of eye ball (d) Older age
21. Human eye is most sensitive to visible light of the wavelength  
 (a) 6050 Å (b) 5500 Å (c) 4500 Å (d) 7500 Å
22. Match the List I with the List II from the combinations shown  
 (I) Presbiopia (A) Sphero-cylindrical lens  
 (II) Hypermetropia (B) Convex lens of proper power may be used close to the eye

- (III) Astigmatism (C) Concave lens of suitable focal length  
 (IV) Myopia (D) Convex spectacle lens of suitable focal length  
 (a) I-A; II-C; III-B; IV-D (b) I-B; II-D; III-C; IV-A (c) I-D; II-B; III-A; IV-C (d) I-D; II-A; III-C; IV-B
23. The human eye has a lens which has a  
 (a) Soft portion at its centre (b) Hard surface  
 (c) Varying refractive index (d) Constant refractive index
24. A man with defective eyes cannot see distinctly object at the distance more than 60 cm from his eyes. The power of the lens to be used will be  
 (a) + 60D (b) - 60D (c) - 1.66D (d)  $\frac{1}{1.66} D$
25. A person's near point is 50 cm and his far point is 3 m. Power of the lenses he requires for  
 (i) Reading and (ii) For seeing distant stars are  
 (a) - 2D and 0.33D (b) 2D and - 0.33D (c) - 2D and 3D (d) 2D and - 3D
26. The focal length of a simple convex lens used as a magnifier is 10 cm. For the image to be formed at a distance of distinct vision ( $D = 25$  cm), the object must be placed away from the lens at a distance of  
 (a) 5 cm (b) 7.14 cm (c) 7.20 cm (d) 16.16 cm
27. A person is suffering from myopic defect. He is able to see clear objects placed at 15 cm. What type and of what focal length of lens he should use to see clearly the object placed 60 cm away  
 (a) Concave lens of 20 cm focal length (b) Convex lens of 20 cm focal length  
 (c) Concave lens of 12 cm focal length (d) Convex lens of 12 cm focal length
28. A person can see a thing clearly when it is at a distance of 1 metre only. If he wishes to see a distance star, he needs a lens of focal length  
 (a) +100 cm (b) - 100 cm (c) +50 cm (d) -50 cm
29. A man suffering from myopia can read a book placed at 10 cm distance. For reading the book at a distance of 60 cm with relaxed vision, focal length of the lens required will be  
 (a) 45 cm (b) - 20 cm (c) -12 cm (d) 30 cm
30. A person can see clearly objects at 100 cm distance. If he wants to see objects at 40 cm distance, then the power of the lens he shall require is  
 (a) +1.5 D (b) - 1.5 D (c) +3.0 D (d) -3.0 D
31. If the distance of the far point for a myopia patient is doubled, the focal length of the lens required to cure it will become  
 (a) Half (b) Double  
 (c) The same but a convex lens (d) The same but a concave lens
32. Image is formed for the short sighted person at  
 (a) Retina (b) Before retina (c) Behind the retina (d) Image is not formed at all
33. A man who cannot see clearly beyond 5 m wants to see stars clearly. He should use a lens of focal length  
 (a) - 100 metre (b) + 5 metre (c) - 5 metre (d) Very large
34. Far point of myopic eye is 250 cm, then the focal length of the lens to be used will be  
 (a) + 250 cm (b) - 250 cm (c) + 250/9 cm (d) - 250/9 cm
35. One can take pictures of objects which are completely invisible to the eye using camera film which are invisible to  
 (a) Ultra-violet rays (b) Sodium light (c) Visible light (d) Infra-red rays

36. In human eye the focussing is done by  
 (a) To and fro movement of eye lens (b) To and fro movement of the retina  
 (c) Change in the convexity of the lens surface (d) Change in the refractive index of the eye fluids
37. The minimum light intensity that can be perceived by the eye is about  $10^{-10}$  watt / metre<sup>2</sup>. The number of photons of wavelength  $5.6 \times 10^{-7}$  metre that must enter per second the pupil of area  $10^{-4}$  metre<sup>2</sup> for vision, is approximately equal to ( $h = 6.6 \times 10^{-34}$  joule - sec)  
 (a)  $3 \times 10^2$  photons (b)  $3 \times 10^6$  photons (c)  $3 \times 10^4$  photons (d)  $3 \times 10^5$  photons
38. A far sighted man who has lost his spectacles, reads a book by looking through a small hole (3-4 mm) in a sheet of paper. The reason will be  
 (a) Because the hole produces an image of the letters at a longer distance  
 (b) Because in doing so, the focal length of the eye lens is effectively increased  
 (c) Because in doing so, the focal length of the eye lens is effectively decreased  
 (d) None of these
39. The maximum focal length of the eye-lens of a person is greater than its distance from the retina. The eye is  
 (a) Always strained in looking at an object (b) Strained for objects at large distances only  
 (c) Strained for objects at short distances only (d) Unstrained for all distances
40. The focal length of a normal eye-lens is about  
 (a) 1 mm (b) 2 cm (c) 25 cm (d) 1
41. The distance of the eye-lens from the retina is  $x$ . For normal eye, the maximum focal length of the eye-lens is  
 (a)  $= x$  (b)  $< x$  (c)  $> x$  (d)  $= 2x$
42. A man wearing glasses of focal length  $+1m$  can clearly see beyond  $1m$   
 (a) If he is farsighted (b) If he is nearsighted (c) If his vision is normal (d) In each of these cases
43. The near point of a person is 50 cm and the far point is 1.5 m. The spectacles required for reading purpose and for seeing distance are respectively  
 (a)  $+2D, -\left(\frac{2}{3}\right)D$  (b)  $+\left(\frac{2}{3}\right)D - 2D$  (c)  $-2D, +\left(\frac{2}{3}\right)D$  (d)  $-\left(\frac{2}{3}\right)D + 2D$
44. A man, wearing glasses of power  $+2D$  can read clearly a book placed at a distance of 40 cm from the eye. The power of the lens required so that he can read at 25 cm from the eye is  
 (a)  $+4.5 D$  (b)  $+4.0 D$  (c)  $+3.5 D$  (d)  $+3.0 D$
45. A person can see clearly between 1 m and 2m. His corrective lenses should be  
 (a) Bifocals with power  $-0.5D$  and additional  $+3.5D$  (b) Bifocals with power  $-1.0D$  and additional  $+3.0 D$   
 (c) Concave with power  $1.0 D$  (d) Convex with power  $0.5 D$
46. While reading the book a man keeps the page at a distance of 2.5 cm from his eye. He wants to read the book by holding the page at 25 cm. What is the nature of spectacles one should advice him to use to completely cure his eye sight  
 (a) Convex lens of focal length 25 cm (b) Concave lens of focal length 25 cm  
 (c) Convex lens of focal length 2.5 cm (d) Concave lens of focal length 2.5 cm
47. The blades of a rotating fan can not be distinguished from each other due to  
 (a) Parallax (b) Power of accommodation (c) Persistence of vision (d) Binocular vision

48. Aperture of the human eye is 2 mm. Assuming the mean wavelength of light to be 5000 Å, the angular resolution limit of the eye is nearly  
 (a) 2 minutes (b) 1 minute (c) 0.5 minute (d) 1.5 minutes
49. If there had been one eye of the man, then  
 (a) Image of the object would have been inverted (b) Visible region would have decreased  
 (c) Image would have not been seen three dimensional (d) (b) and (c) both
50. A man can see the object between 15cm and 30cm. He uses the lens to see the far objects. Then due to the lens used, the near point will be at  
 (a)  $\frac{10}{3}$  cm (b) 30 cm (c) 15 cm (d)  $\frac{100}{3}$  cm
51. A presbyopic patient has near point as 30 cm and far point as 40 cm. The dioptric power for the corrective lens for seeing distant objects is  
 (a) 40 D (b) 4 D (c) 2.5 D (d) 0.25 D
52. A man swimming under clear water is unable to see clearly because  
 (a) The size of the aperture decreases (b) The size of the aperture increases  
 (c) The focal length of eye lens increases (d) The focal length of eye lens decreases
53. The distance between retina and eye-lens in a normal eye is 2.0 cm. The accommodated power of eye lens range from  
 (a) 45 D to 50 D (b) 50 D to 54 D (c) 10 D to 16 D (d) 5 D to 8 D
54. If the eye is taken as a spherical ball of radius 1 cm, the range of accommodated focal length of eye-lens is  
 (a) 1.85 cm to 2.0 cm (b) 1.0 cm to 2.8 cm (c) 1.56 cm to 2.5 cm (d) 1.6 cm to 2.0 cm
55. A person cannot read printed matter within 100 cm from his eye. The power of the correcting lens required to read at 20 cm from his eye if the distance between the eye lens and the correcting lens is 2 cm is  
 (a) 4.8 D (b) 1.25 D (c) 4.25 D (d) 4.55 D
56. A student having  $-1.5$  D spectacles uses a lens of focal length 5 cm as a simple microscope to read minute scale divisions in the laboratory. The least distance of distinct vision without glasses is 20 cm for the student. The maximum magnifying power he gets with spectacles on is  
 (a) 6 (b) 9 (c) 5 (d) 4

**Microscope**

57. In a compound microscope the object of  $f_o$  and eyepiece of  $f_e$  are placed at distance  $L$  such that  $L$  equals  
 (a)  $f_o + f_e$  (b)  $f_o - f_e$   
 (c) Much greater than  $f_o$  or  $f_e$  (d) Need not depend either value of focal lengths
58. In a simple microscope, if the final image is located at infinity then its magnifying power is  
 (a)  $\frac{25}{f}$  (b)  $\frac{D}{25}$  (c)  $\frac{f}{25}$  (d)  $\frac{f}{D^{+1}}$
59. In a simple microscope, if the final image is located at 25 cm from the eye placed close to the lens, then the magnifying power is

- (a)  $\frac{25}{f}$  (b)  $1 + \frac{25}{f}$  (c)  $\frac{f}{25}$  (d)  $\frac{f}{25} + 1$
60. The maximum magnification that can be obtained with a convex lens of focal length 2.5 cm is (the least distance of distinct vision is 25 cm)
- (a) 10 (b) 0.1 (c) 62.5 (d) 11
61. In a compound microscope, the intermediate image is
- (a) Virtual, erect and magnified (b) Real, erect and magnified  
(c) Real, inverted and magnified (d) Virtual, erect and reduced
62. A compound microscope has two lenses. The magnifying power of one is 5 and the combined magnifying power is 100. The magnifying power of the other lens is
- (a) 10 (b) 20 (c) 50 (d) 25
63. Wavelength of light used in an optical instrument are  $\lambda_1 = 4000 \text{ \AA}$  and  $\lambda_2 = 5000 \text{ \AA}$ , then ratio of their respective resolving power (corresponding to  $\lambda_1$  and  $\lambda_2$ ) is
- (a) 16 : 25 (b) 9 : 1 (c) 4 : 5 (d) 5 : 4
64. The angular magnification of a simple microscope can be increased by increasing
- (a) Focal length of lens (b) Size of object (c) Aperture of lens (d) Power of lens
65. The magnification produced by the objective lens and the eye lens of a compound microscope are 25 and 6 respectively. The magnifying power of this microscope is
- (a) 19 (b) 31 (c) 150 (d)  $\sqrt{150}$
66. The length of the compound microscope is 14 cm. The magnifying power for relaxed eye is 25. If the focal length of eye lens is 5 cm, then the object distance for objective lens will be
- (a) 1.8 cm (b) 1.5 cm (c) 2.1 cm (d) 2.4 cm
67. The magnifying power of a simple microscope is 6. The focal length of its lens in metres will be, if least distance of distinct vision is 25 cm
- (a) 0.05 (b) 0.06 (c) 0.25 (d) 0.12
68. Relative difference of focal lengths of objective and eye lens in the microscope and telescope is given as
- (a) It is equal in both (b) It is more in telescope (c) It is more in microscope (d) It may be more in any one
69. Three objective focal lengths ( $f_o$ ) and two eye piece focal lengths ( $f_e$ ) are available for a compound microscope. By combining these two, the magnification of microscope will be maximum when
- (a)  $f_o = f_e$  (b)  $f_o \gg f_e$  (c)  $f_o$  and  $f_e$  both are small (d)  $f_o \gg f_e$
70. If the red light is replaced by blue light illuminating the object in a microscope the resolving power of the microscope
- (a) Decreases (b) Increases (c) Gets halved (d) Remains unchanged
71. In case of a simple microscope, the object is placed at
- (a) Focus  $f$  of the convex lens (b) A position between  $f$  and  $2f$   
(c) Beyond  $2f$  (d) Between the lens and  $f$
72. In a compound microscope cross-wires are fixed at the point
- (a) Where the image is formed by the objective (b) Where the image is formed by the eye-piece

- (c) Where the focal point of the objective lies (d) Where the focal point of the eye-piece lies
73. The length of the tube of a microscope is 10 cm. The focal lengths of the objective and eye lenses are 0.5 cm and 1.0 cm. The magnifying power of the microscope is about  
 (a) 5 (b) 23 (c) 166 (d) 500
74. Least distance of distinct vision is 25 cm. Magnifying power of simple microscope of focal length 5 cm is  
 (a) 1/5 (b) 5 (c) 1/6 (d) 6
75. The objective of a compound microscope is essentially  
 (a) A concave lens of small focal length and small aperture (b) Convex lens of small focal length and large aperture  
 (c) Convex lens of large focal length and large aperture (d) Convex lens of small focal length and small aperture
76. For relaxed eye, the magnifying power of a microscope is  
 (a)  $-\frac{v_o}{u_o} \times \frac{D}{f_e}$  (b)  $-\frac{v_o}{u_o} \times \frac{f_e}{D}$  (c)  $\frac{u_o}{v_o} \times \frac{D}{f_e}$  (d)  $\frac{u_o}{v_o} \times \left(-\frac{D}{f_e}\right)$
77. A person using a lens as a simple microscope sees an  
 (a) Inverted virtual image (b) Inverted real magnified image  
 (c) Upright virtual image (d) Upright real magnified image
78. The focal length of the objective lens of a compound microscope is  
 (a) Equal to the focal length of its eye piece (b) Less than the focal length of eye piece  
 (c) Greater than the focal length of eye piece (d) Any of the above three
79. To produce magnified erect image of a far object, we will be required along with a convex lens, is  
 (a) Another convex lens (b) Concave lens (c) A plane mirror (d) A concave mirror
80. An object placed 10 cm in front of a lens has an image 20 cm behind the lens. What is the power of the lens (in dioptries)  
 (a) 1.5 (b) 3.0 (c) - 15.0 (d) +15.0
81. Resolving power of a microscope depends upon  
 (a) The focal length and aperture of the eye lens (b) The focal lengths of the objective and the eye lens  
 (c) The apertures of the objective and the eye lens (d) The wavelength of light illuminating the object
82. If the focal length of the objective lens is increased then  
 (a) Magnifying power of microscope will increase but that of telescope will decrease  
 (b) Magnifying power of microscope and telescope both will increase  
 (c) Magnifying power of microscope and telescope both will decrease  
 (d) Magnifying power of microscope will decrease but that of telescope will increase

83. If in compound microscope  $m_1$  and  $m_2$  be the linear magnification of the objective lens and eye lens respectively, then magnifying power of the compound microscope will be  
 (a)  $m_1 - m_2$  (b)  $\sqrt{m_1 + m_2}$  (c)  $(m_1 + m_2)/2$  (d)  $m_1 \times m_2$
84. The magnifying power of a microscope with an objective of 5 mm focal length is 400. The length of its tube is 20 cm. Then the focal length of the eye-piece is  
 (a) 200 cm (b) 160 cm (c) 2.5 cm (d) 0.1 cm
85. In a compound microscope, if the objective produces an image  $I_o$  and the eye piece produces an image  $I_e$ , then  
 (a)  $I_o$  is virtual but  $I_e$  is real (b)  $I_o$  is real but  $I_e$  is virtual (c)  $I_o$  and  $I_e$  are both real (d)  $I_o$  and  $I_e$  are both virtual
86. In an electron microscope if the potential is increased from 20 kV to 80 kV, the resolving power of the microscope will change from  $R$  to  
 (a)  $R/4$  (b)  $4R$  (c)  $2R$  (d)  $R/2$
87. When the length of a microscope tube increases, its magnifying power  
 (a) Decreases (b) Increases (c) Does not change (d) May decrease or increase
88. An electron microscope is superior to an optical microscope in  
 (a) Having better resolving power (b) Being easy to handle  
 (c) Low cost (d) Quickness of observation
89. In a compound microscope magnification will be large, if the focal length of the eye piece is  
 (a) Large (b) Smaller (c) Equal to that of objective (d) Less than that of objective
90. An electron microscope gives better resolution than optical microscope because  
 (a) Electrons are abundant (b) Electrons can be focused nicely  
 (c) Effective wavelength of electron is small (d) None of these
91. A man is looking at a small object placed at his near point. Without altering the position of his eye or the object, he puts a simple microscope of magnifying power 5X before his eyes. The angular magnification achieved is  
 (a) 5 (b) 2.5 (c) 1 (d) 0.2
92. The focal length of the objective of a compound microscope is  $f_0$  and its distance from the eyepiece is  $L$ . The object is placed at a distance  $u$  from the objective. For proper working of the instrument  
 (a)  $L < u$  (b)  $L > u$  (c)  $f_0 < L < 2f_0$  (d)  $L > 2f_0$
93. Find the maximum magnifying power of a compound microscope having a 25 diopter lens as the objective, a 5 diopter lens as the eyepiece and the separation 30 cm between the two lenses. The least distance for clear vision is 25 cm  
 (a) 8.4 (b) 7.4 (c) 9.4 (d) 10.4
94. The focal length of the objective and the eye-piece of a microscope are 2 cm and 5 cm respectively and the distance between them is 30 cm. If the image seen by the eye is 25 cm from the eye-piece, the distance of the object from the objective is  
 (a) 0.8 cm (b) 2.3 cm (c) 0.4 cm (d) 1.2 cm



95. The focal length of objective and eye-piece of a microscope are 1 cm and 5 cm respectively. If the magnifying power for relaxed eye is 45, then length of the tube is  
 (a) 6 cm (b) 9 cm (c) 12 cm (d) 15 cm
96. A microscope has an objective of focal length 1.5 cm and an eye-piece of focal length 2.5 cm. If the distance between objective and eye-piece is 25 cm. What is the approximate value of magnification produced for relaxed eye is  
 (a) 75 (b) 110 (c) 140 (d) 25
97. The magnifying power of a microscope is generally marked as 10X, 100 X, etc. These markings are for a normal relaxed eye. A microscope marked 10X is used by an old man having his near point at 40 cm. The magnifying power of the microscope for the old man with his eyes completely relaxed is  
 (a) 10 (b) 18 (c) 12 (d) 16
98. If the focal length of objective and eye lens are 1.2 cm and 3 cm respectively and the object is put 1.25 cm away from the objective lens and the final image is formed at infinity. The magnifying power of the microscope is  
 (a) 150 (b) 200 (c) 250 (d) 400
99. A compound microscope is adjusted for viewing the distant image of an object, the distance of the object from the object glass is now slightly increased, what re-adjustment of the instrument would be necessary for obtaining a distant image again  
 (a) Objective should be moved away from the eye-piece (b) Eye-piece should be moved towards the objective  
 (c) Both should be moved towards each other (d) Both should be moved away from each other
100. When the object is self-luminous, the resolving power of a microscope is given by the expression  
 (a)  $\frac{2\mu \sin \theta}{\lambda}$  (b)  $\frac{\mu \sin \theta}{\lambda}$  (c)  $\frac{2\mu \cos \theta}{\lambda}$  (d)  $\frac{2\mu}{\lambda}$
101. In a compound microscope, maximum magnification is obtained when the final image  
 (a) Is formed at infinity (b) Is formed at the least of distinct vision  
 (c) Coincides with the object (d) Coincides with the objective lens
102. How should people wearing spectacles work with a microscope  
 (a) They should keep on wearing their spectacles  
 (b) They should take off their spectacles  
 (c) They may keep on wearing or take off their spectacles, It makes no difference  
 (d) They cannot use a microscope at all

**Telescope**

103. The focal length of the objective and eyepiece of an astronomical telescope for normal adjustments are 50 cm and 5 cm. The length of the telescope should be  
 (a) 50 cm (b) 55 cm (c) 60 cm (d) 45 cm
104. The resolving power of an astronomical telescope is 0.2 seconds. If the central half portion of the objective lens is covered, the resolving power will be  
 (a) 0.1 sec (b) 0.2 sec (c) 1.0 sec (d) 0.6 sec

105. If  $F_o$  and  $F_e$  are the focal length of the objective and eye-piece respectively of a telescope, then its magnifying power will be
- (a)  $F_o + F_e$                       (b)  $F_o \times F_e$                       (c)  $F_o / F_e$                       (d)  $\frac{1}{2}(F_o + F_e)$
106. The length of an astronomical telescope for normal vision (relaxed eye) ( $f_o$  = focal length of objective lens and  $f_e$  = focal length of eye lens) is
- (a)  $f_o \times f_e$                       (b)  $\frac{f_o}{f_e}$                       (c)  $f_o + f_e$                       (d)  $f_o - f_e$
107. A telescope of diameter  $2m$  uses light of wavelength  $5000 \text{ \AA}$  for viewing stars. The minimum angular separation between two stars whose image is just resolved by this telescope is
- (a)  $4 \times 10^{-4} \text{ rad}$                       (b)  $0.25 \times 10^{-6} \text{ rad}$                       (c)  $0.31 \times 10^{-6} \text{ rad}$                       (d)  $5.0 \times 10^{-3} \text{ rad}$
108. The aperture of the objective lens of a telescope is made large so as to
- (a) Increase the magnifying power of the telescope                      (b) Increase the resolving power of the telescope  
(c) Make image aberration less                      (d) Focus on distant objects
109. The distance of the moon from earth is  $3.8 \times 10^5 \text{ km}$ . The eye is most sensitive to light of wavelength  $5500 \text{ \AA}$ . The separation of two points on the moon that can be resolved by a  $500 \text{ cm}$  telescope will be
- (a)  $51 \text{ m}$                       (b)  $60 \text{ m}$                       (c)  $70 \text{ m}$                       (d) All of the above
110. To increase both the resolving power and magnifying power of a telescope
- (a) Both the focal length and aperture of the objective has to be increased  
(b) The focal length of the objective has to be increased  
(c) The aperture of the objective has to be increased  
(d) The wavelength of light has to be decreased
111. The focal lengths of the objective and eye lenses of a telescope are respectively  $200 \text{ cm}$  and  $5 \text{ cm}$ . The maximum magnifying power of the telescope will be
- (a)  $-40$                       (b)  $-48$                       (c)  $-60$                       (d)  $-100$
112. A telescope has an objective of focal length  $50 \text{ cm}$  and an eye piece of focal length  $5 \text{ cm}$ . The least distance of distinct vision is  $25 \text{ cm}$ . The telescope is focussed for distinct vision on a scale  $200 \text{ cm}$  away. The separation between the objective and the eye-piece is
- (a)  $75 \text{ cm}$                       (b)  $60 \text{ cm}$                       (c)  $71 \text{ cm}$                       (d)  $74 \text{ cm}$
113. In a laboratory four convex lenses  $L_1, L_2, L_3$  and  $L_4$  of focal lengths  $2, 4, 6$  and  $8 \text{ cm}$  respectively are available. Two of these lenses form a telescope of length  $10 \text{ cm}$  and magnifying power  $4$ . The objective and eye lenses are
- (a)  $L_2, L_3$                       (b)  $L_1, L_4$                       (c)  $L_3, L_2$                       (d)  $L_4, L_1$
114. Four lenses of focal length  $+15 \text{ cm}$ ,  $+20 \text{ cm}$ ,  $+150 \text{ cm}$  and  $+250 \text{ cm}$  are available for making an astronomical telescope. To produce the largest magnification, the focal length of the eye-piece should be
- (a)  $+15 \text{ cm}$                       (b)  $+20 \text{ cm}$                       (c)  $+150 \text{ cm}$                       (d)  $+250 \text{ cm}$
115. In a terrestrial telescope, the focal length of objective is  $90 \text{ cm}$ , of inverting lens is  $5 \text{ cm}$  and of eye lens is  $6 \text{ cm}$ . If the final image is at  $30 \text{ cm}$ , then the magnification will be
- (a)  $21$                       (b)  $12$                       (c)  $18$                       (d)  $15$

116. The focal lengths of the objective and the eyepiece of an astronomical telescope are 20 cm and 5 cm respectively. If the final image is formed at a distance of 30 cm from the eye piece, find the separation between the lenses for distinct vision
- (a) 32.4 cm                      (b) 42.3 cm                      (c) 24.3 cm                      (d) 30.24 cm
117. Resolving power of reflecting type telescope increases with
- (a) Decrease in wavelength of incident light                      (b) Increase in wavelength of incident light  
(c) Increase in diameter of objective lens                      (d) None of these
118. A planet is observed by an astronomical refracting telescope having an objective of focal length 16 m and an eye-piece of focal length 2 cm [IIT-JEE 1992; Roorkee 2000]
- (a) The distance between the objective and the eye-piece is 16.02 m  
(b) The angular magnification of the planet is 800  
(c) The image of the planet is inverted  
(d) All of the above
119. The astronomical telescope consists of objective and eye-piece. The focal length of the objective is
- (a) Equal to that of the eye-piece                      (b) Greater than that of the eye-piece  
(c) Shorter than that of the eye-piece                      (d) Five times shorter than that of the eye-piece
120. The diameter of the objective of a telescope is  $a$ , the magnifying power is  $m$  and wavelength of light is  $\lambda$ . The resolving power of the telescope is
- (a)  $(1.22\lambda)/a$                       (b)  $(1.22a)/\lambda$                       (c)  $\lambda m/(1.22a)$                       (d)  $a/(1.22\lambda m)$
121. An astronomical telescope has an angular magnification of magnitude 5 for distant objects. The separation between the objective and the eyepiece is 36 cm and final image is formed at infinity. The focal lengths of the objective and eyepiece are respectively [IIT-JEE 1989]
- (a) 20 cm, 16 cm                      (b) 50 cm, 10 cm                      (c) 30 cm, 6 cm                      (d) 45 cm, -9 cm
122. A photograph of the moon was taken with telescope. Later on, it was found that a housefly was sitting on the objective lens of the telescope. In photograph
- (a) The image of housefly will be reduced                      (b) There is a reduction in the intensity of the image  
(c) There is an increase in the intensity of the image                      (d) The image of the housefly will be enlarged
123. The magnifying power of a telescope is  $M$ . If the focal length of eye piece is doubled, then the magnifying power will become
- (a)  $2M$                       (b)  $M/2$                       (c)  $\sqrt{2M}$                       (d)  $3M$
124. The minimum magnifying power of a telescope is  $M$ . If the focal length of its eyelens is halved, the magnifying power will become
- (a)  $M/2$                       (b)  $2M$                       (c)  $3M$                       (d)  $4M$
125. The final image in an astronomical telescope is
- (a) Real and erect                      (b) Virtual and inverted                      (c) Real and inverted                      (d) Virtual and erect
126. The astronomical telescope has two lenses of focal powers 0.5 D and 20 D. Its magnifying power will be
- (a) 40                      (b) 10                      (c) 100                      (d) 35
127. An astronomical telescope of ten-fold angular magnification has a length of 44 cm. The focal length of the objective is
- (a) 4 cm                      (b) 40 cm                      (c) 44 cm                      (d) 440 cm

128. A telescope consisting of an objective of focal length 100 cm and a single eye lens of focal length 10 cm is focussed on a distant object in such a way that parallel rays emerge from the eye lens. If the object subtends an angle of  $2^\circ$  at the objective, the angular width of the image is  
 (a)  $20^\circ$  (b)  $1/6^\circ$  (c)  $10^\circ$  (d)  $24^\circ$
129. When diameter of the aperture of the objective of an astronomical telescope is increased, its  
 (a) Magnifying power is increased and resolving power is decreased  
 (b) Magnifying power and resolving power both are increased  
 (c) Magnifying power remains the same but resolving power is increased  
 (d) Magnifying power and resolving power both are decreased
130. The focal length of objective and eye-piece of a telescope are 100 cm and 5 cm respectively. Final image is formed at least distance of distinct vision. The magnification of telescope is  
 (a) 20 (b) 24 (c) 30 (d) 36
131. A simple telescope, consisting of an objective of focal length 60 cm and single eye lens of focal length 5 cm is focussed on a distant object in such a way that parallel rays come out from the eye lens. If the object subtends an angle  $2^\circ$  at the objective, the angular width of the image  
 (a)  $10^\circ$  (b)  $24^\circ$  (c)  $50^\circ$  (d)  $1/6^\circ$
132. The diameter of the objective of the telescope is 0.1 metre and wavelength of light is  $6000 \text{ \AA}$ . Its resolving power would be approximately  
 (a)  $7.32 \times 10^{-6} \text{ radian}$  (b)  $1.36 \times 10^6 \text{ radian}$  (c)  $7.32 \times 10^{-5} \text{ radian}$  (d)  $1.36 \times 10^5 \text{ radian}$
133. A Galilean telescope has objective and eye-piece of focal lengths 200 cm and 2 cm respectively. The magnifying power of the telescope for normal vision is  
 (a) 90 (b) 100 (c) 108 (d) 198
134. All of the following statements are correct except  
 (a) The total focal length of an astronomical telescope is the sum of the focal lengths of its two lenses  
 (b) The image formed by the astronomical telescope is always erect because the effect of the combination of the two lenses is divergent  
 (c) The magnification of an astronomical telescope can be increased by decreasing the focal length of the eye-piece  
 (d) The magnifying power of the refracting type of astronomical telescope is the ratio of the focal length of the objective to that of the eye-piece
135. The length of a telescope is 36 cm. The focal length of its lenses can be  
 (a) 30 cm, 6 cm (b) - 30 cm, - 6 cm (c) - 30 cm, - 6 cm (d) - 30 cm, 6 cm
136. The diameter of the objective lens of telescope is 5.0 m and wavelength of light is  $6000 \text{ \AA}$ . The limit of resolution of this telescope will be  
 (a) 0.03 sec (b) 3.03 sec (c) 0.06 sec (d) 0.15 sec
137. If tube length of astronomical telescope is 105 cm and magnifying power is 20 for normal setting, calculate the focal length of objective  
 (a) 100 cm (b) 10 cm (c) 20 cm (d) 25 cm
138. Radio telescope is used to see  
 (a) Distant stars and planets (b) Sun and to measure its temperature  
 (c) Stars and to measure diameters (d) None of these

139. Four lenses with focal length  $\pm 15\text{ cm}$  and  $\pm 150\text{ cm}$  are being placed for use as a telescopic objective. The focal length of the lens which produces the largest magnification with a given eye-piece is  
 (a)  $-15\text{ cm}$  (b)  $+150\text{ cm}$  (c)  $-150\text{ cm}$  (d)  $+15\text{ cm}$
140. The image of a star (effectively a point source) is made by a convergent lens of focal length  $50\text{ cm}$  and diameter of aperture  $5.0\text{ cm}$ . If the lens is ideal, and the effective wavelength in image formation is taken as  $5 \times 10^{-5}\text{ cm}$ , the diameter of the image formed will be nearest to  
 (a) Zero (b)  $10^{-6}\text{ cm}$  (c)  $10^{-5}\text{ cm}$  (d)  $10^{-3}\text{ cm}$
141. To increase the magnifying power of a telescope ( $f_o$  = focal length of the objective and  $f_e$  = focal length of the eye lens)  
 (a)  $f_o$  should be large and  $f_e$  should be small (b)  $f_o$  should be small and  $f_e$  should be large  
 (c)  $f_o$  and  $f_e$  both should be large (d)  $f_o$  and  $f_e$  both should be small
142. The limit of resolution of a  $100\text{ cm}$  telescope ( $\lambda = 5.5 \times 10^{-7}\text{ m}$ ) is [ ]  
 (a)  $0.14''$  (b)  $0.3''$  (c)  $1'$  (d)  $1''$
143. In a reflecting astronomical telescope, if the objective (a spherical mirror) is replaced by a parabolic mirror of the same focal length and aperture, then [IIT-JEE 1993]  
 (a) The final image will be erect (b) The larger image will be obtained  
 (c) The telescope will gather more light (d) Spherical aberration will be absent
144. A planet is observed by an astronomical refracting telescope having an objective of focal length  $16\text{ m}$  and an eyepiece of focal length  $2\text{ cm}$  [IIT-JEE 1993]  
 (a) The distance between the objective and the eyepiece is  $16.02\text{ m}$   
 (b) The angular magnification of the planet is 800  
 (c) The image of the planet is inverted  
 (d) The objective is larger than the eyepiece
145. The average distance between the earth and moon is  $38.6 \times 10^4\text{ km}$ . The minimum separation between the two points on the surface of the moon that can be resolved by a telescope whose objective lens has a diameter of  $5\text{ m}$  with  $\lambda = 6000\text{ \AA}$  is  
 (a)  $5.65\text{ m}$  (b)  $28.25\text{ m}$  (c)  $11.30\text{ m}$  (d)  $56.51\text{ m}$
146. The focal length of the objective and eye piece of a telescope are respectively  $60\text{ cm}$  and  $10\text{ cm}$ . The magnitude of the magnifying power when the image is formed at infinity is  
 (a) 50 (b) 6 (c) 70 (d) 5
147. The focal length of an objective of a telescope is  $3\text{ metre}$  and diameter  $15\text{ cm}$ . Assuming for a normal eye, the diameter of the pupil is  $3\text{ mm}$  for its complete use, the focal length of the eye piece must be  
 (a)  $6\text{ cm}$  (b)  $6.3\text{ cm}$  (c)  $20\text{ cm}$  (d)  $60\text{ cm}$
148. An opera glass (Gallilean telescope) measures  $9\text{ cm}$  from the objective to the eyepiece. The focal length of the objective is  $15\text{ cm}$ . Its magnifying power is  
 (a) 2.5 (b)  $2/5$  (c)  $5/3$  (d) 0.4
149. The focal length of the objective and eye lens of an astronomical telescope are respectively  $2\text{ m}$  and  $5\text{ cm}$ . The final image is formed at (i) least distance of distinct vision (ii) infinity. The magnifying power in both cases will be  
 (a)  $-48, -40$  (b)  $-40, -48$  (c)  $-40, 48$  (d)  $-48, 40$
150. An optical device that enables an observer to see over or around opaque objects, is called  
 (a) Microscope (b) Telescope (c) Periscope (d) Hydrometer

151. The magnifying power of a telescope can be increased by  
 (a) Increasing focal length of the system (b) Fitting eye piece of high power  
 (c) Fitting eye piece of low power (d) Increasing the distance of objects
152. An achromatic telescope objective is to be made by combining the lenses of flint and crown glasses. This proper choice is  
 (a) Convergent of crown and divergent of flint (b) Divergent of crown and convergent of flint  
 (c) Both divergent (d) Both convergent
153. An observer looks at a tree of height 15 m with a telescope of magnifying power 10. To him, the tree appears  
 (a) 10 times taller (b) 15 times taller (c) 10 times nearer (d) 15 times nearer
154. The magnification produced by an astronomical telescope for normal adjustment is 10 and the length of the telescope is 1.1 m. The magnification when the image is formed at least distance of distinct vision ( $D = 25 \text{ cm}$ ) is  
 (a) 14 (b) 6 (c) 16 (d) 18
155. The objective of a telescope has a focal length of 1.2 m. it is used to view a 10.0 m tall tower 2 km away. What is the height of the image of the tower formed by the objective  
 (a) 2 mm (b) 4 mm (c) 6 mm (d) 8 mm
156. A giant telescope in an observatory has an objective of focal length 19 m and an eye-piece of focal length 1.0 cm. In normal adjustment, the telescope is used to view the moon. What is the diameter of the image of the moon formed by the objective? The diameter of the moon is  $3.5 \times 10^6 \text{ m}$  and the radius of the lunar orbit round the earth is  $3.8 \times 10^8 \text{ m}$   
 (a) 10 cm (b) 12.5 cm (c) 15 cm (d) 17.5 cm
157. The aperture of the largest telescope in the world is  $\approx 5 \text{ metre}$ . If the separation between the moon and the earth is  $\approx 4 \times 10^5 \text{ km}$  and the wavelength of the visible light is  $\approx 5000 \text{ \AA}$ , then the minimum separation between the objects on the surface of the moon which can be just resolved is  
 (a) 1 metre approximately (b) 10 metre approximately (c) 50 metre approximately (d) 200 metre approximately
158. In Galileo's telescope, magnifying power for normal vision is 20 and power of eye-piece is  $-20 \text{ D}$ . Distance between the objective and eye-piece should be  
 (a) 90 cm (b) 95 cm (c) 100 cm (d) 105 cm
159. The least resolve angle by a telescope using objective of aperture 5 m and light of wavelength = 4000 A.U. is nearly  
 (a)  $\frac{1}{50}^\circ$  (b)  $\frac{1}{50} \text{ sec}$  (c)  $\frac{1}{50} \text{ minute}$  (d)  $\frac{1}{500} \text{ sec}$
160. The limit of resolution of a 10 cm telescope for visible light of wavelength 6000 Å is approximately  
 (a) 0.1 s or arc (b)  $30^\circ$  (c)  $\left(\frac{1}{6}\right)^\circ$  (d) None of these
161. An eye-piece of a telescope with a magnification of 100 has a power of 20 diopters. The object of this telescope has a power of  
 (a) 2 diopters (b) 0.2 diopters (c) 2000 diopters (d) 20 diopters
162. The Yerkes Observatory telescope has a large telescope with objective of diameter of about 1 m. Assuming wavelength of light to be  $6 \times 10^{-7} \text{ m}$ , the angular distance  $\theta$  between two stars which can just be resolved is  
 (a)  $(7.3 \times 10^{-7})^\circ$  (b)  $7.3 \times 10^{-7} \text{ rad}$  (c)  $\frac{1}{40}$  of a second (d) None of these
163. A Galilean telescope measures 9 cm from the objective to the eye-piece. The focal length of the objective is 15 cm. Its magnifying power is  
 (a) 2.5 (b) 2/5 (c) 5/3 (d) 0.4

164. For seeing a cricket match, we prefer binoculars to the terrestrial telescope, because
- (a) Binoculars give three-dimensional view (b) Terrestrial telescope gives inverted image  
(c) To avoid chromatic aberration (d) To have larger magnification
165. A simple two lens telescope has an objective of focal length 50 cm and an eye-piece of 2.5 cm. The telescope is pointed at an object at a very large distance which subtends at an angle of 1 milliradian on the naked eye. The eye piece is adjusted so that the final virtual image is formed at infinity. The size of the real image formed by the objective is
- (a) 5 mm (b) 1 mm (c) 0.5 mm (d) 0.1 mm
166. The objective of a telescope, after focussing for infinity is taken out and a slit of length  $L$  is placed in its position. A sharp image of the slit is formed by the eye-piece at a certain distance from it on the other side. The length of this image is  $l$ , then magnification of telescope is
- (a)  $\frac{l}{2L}$  (b)  $\frac{2L}{l}$  (c)  $\frac{l}{L}$  (d)  $\frac{L}{l}$
167. An astronomical telescope in normal adjustment receives light from a distant source  $S$ . The tube length is now decreased slightly
- (a) A virtual image of  $S$  will be formed at a finite distance  
(b) No image will be formed  
(c) A small, real image of  $S$  will be formed behind the eye-piece, close to it  
(d) A large, real image of  $S$  will be formed behind the eye-piece, far away from it
168. A telescope consisting of object glass of power + 2 D and eye-glass of power + 20 D is focussed on an object 1m from the object glass. The final image is seen with completely relaxed eye. The magnifying power of the telescope is
- (a) 20 (b) 41 (c) 24 (d) 49.2
169. An astronomical telescope and a Galilean telescope use identical objective lenses. They have the same magnification, when both are in normal adjustment. The eye-piece of the astronomical telescope has a focal length  $f$
- (a) The tube lengths of the two telescopes differ by  $f$  (b) The tube lengths of the two telescopes differ by  $2f$   
(c) The Galilean telescope has a shorter tube length (d) The Galilean telescope has a longer tube length